Erratum

Analysis of Splicing In Vitro Using Extracts of *Saccharomyces cerevisiae*

Manuel Ares Jr.

Cold Spring Harb Protoc 2013; doi: 10.1101/pdb.prot078121

The recipe entitled “Native Quench Buffer” was incorrect when this protocol was first published. The correct version of the recipe is below. In addition, the HTML version of the recipe and the PDF version of the protocol have been amended accordingly.

Native Quench Buffer

Reagents	Quantity (for 1 mL)	Final concentration
HEPES-KOH (1 M, pH 7.4)	50 μ L	50 mM
MgOAc (1 M)	2 μ L	2 mM
Total yeast or other RNA (10 mg/mL)	200 μ L	2 mg/mL
H ₂ O	to 1 mL	
Store at -20°C .

Cold Spring Harbor Protocols

Analysis of Splicing In Vitro Using Extracts of *Saccharomyces cerevisiae*

Manuel Ares Jr.

Cold Spring Harb Protoc; doi: 10.1101/pdb.err089888

Email Alerting Service

Receive free email alerts when new articles cite this article - [click here](#).

Subject Categories

Browse articles on similar topics from *Cold Spring Harbor Protocols*.

To subscribe to *Cold Spring Harbor Protocols* go to:
<http://cshprotocols.cshlp.org/subscriptions>