Topic Introduction

Whole-Cell Recording in the Awake Brain

Doyun Lee^{1,3} and Albert K. Lee^{2,3}

¹Center for Cognition and Sociality, Institute for Basic Science, Daejeon 34141, Republic of Korea; ²Howard Hughes Medical Institute, Janelia Research Campus, Ashburn, Virginia 20147

Intracellular recording is an essential technique for investigating cellular mechanisms underlying complex brain functions. Despite the high sensitivity of the technique to mechanical disturbances, intracellular recording has been applied to awake, behaving, and even freely moving, animals. Here we summarize recent advances in these methods and their application to the measurement and manipulation of membrane potential dynamics for understanding neuronal computations in behaving animals.

INTRODUCTION

Intracellular recording has served as a key technique for investigating the cellular and circuit mechanisms underlying neural activity. Because of its high sensitivity to mechanical disturbances, intracellular recording previously was limited primarily to stable preparations such as brain slices and, more recently, the intact brains of anesthetized animals (Pei et al. 1991; Ferster and Jagadeesh 1992; Metherate et al. 1992; Moore and Nelson 1998; Zhu and Connors 1999; Margrie et al. 2002). Thus, it had not been straightforward to link the accumulated knowledge regarding synaptic and intrinsic mechanisms from experiments conducted mostly in vitro with higher brain functions that generally involve active participation of the animal. The development of intracellular recording methods for awake behaving animals across multiple species (Covey et al. 1996; Fee 2000; Aksay et al. 2001; Margrie et al. 2002; Wilson et al. 2004; Lee et al. 2006, 2014; Harvey et al. 2009; Long et al. 2010; English et al. 2014; Tan et al. 2014) has therefore played and will continue to play a critical role in understanding the neural basis of cognitive processes (for a review, see Long and Lee (2012)).

INTRACELLULAR RECORDINGS IN HEAD-FIXED BEHAVING ANIMALS

Information processing in the brain is affected by vigilance or arousal states. Therefore, some recordings need to be performed during wakefulness to fully understand neuronal computations. Several intracellular recording studies have demonstrated differences in subthreshold membrane potential dynamics during anesthetized or sleep versus awake states (Steriade et al. 2001; Mahon et al. 2006; Constantinople and Bruno 2011; Haider et al. 2013). Furthermore, animals do not simply passively receive sensory input; they actively interact with the environment, which also affects information processing. For instance, whole-cell recordings in the barrel cortex of head-fixed mice revealed

³Correspondence: leedoyun@ibs.re.kr; leea@janelia.hhmi.org

From the Ion Channels collection, edited by Paul J. Kammermeier, Ian Duguid, and Stephan Brenowitz.

© 2017 Cold Spring Harbor Laboratory Press

Cite this introduction as *Cold Spring Harb Protoc*; doi:10.1101/pdb.top087304

membrane potential fluctuations synchronized to active whisker movements. Different neurons had different phase offsets, which could contribute to position coding (Crochet and Petersen 2006).

Intracellular recording has also been performed in head-fixed animals engaged in activities involving much larger bodily motions such as walking, running, or wing-beating. It had been shown previously that locomotion affects the firing rate of orientation-tuned visual neurons (Niell and Stryker 2010). Cellular mechanisms for such gain modulation have been investigated using whole-cell recordings in head-fixed mice that alternated between periods of running and sitting still on a spherical treadmill while being presented with visual stimuli (Bennett et al. 2013; Polack et al. 2013). Whole-cell recordings obtained from flies fixed in place at the head but free to move their wings were used to study flight-dependent gain modulation in visual neurons from the lobula plate (Maimon et al. 2010).

Head-fixed preparations that allow intracellular recordings to be performed during locomotion on a treadmill have been combined with visually simulated spatial environments where visual scenes are updated based on the animal's movements. Such virtual reality systems allow the study of the brain's navigation system. Whole-cell recordings from mouse hippocampal CA1 place cells revealed intracellular features such as a ramping subthreshold depolarization of the membrane potential during traversals through place fields and phase precession of the intracellular theta-band oscillation with respect to the extracellular local field potential theta rhythm, providing deeper understanding into place cell mechanisms (Harvey et al. 2009). Similar approaches have been used for measuring intracellular features of entorhinal grid cells, allowing investigation into the mechanisms underlying their periodic firing pattern as a function of the animal's location in space (Domnisoru et al. 2013; Schmidt-Hieber and Häusser 2013), as well as for assessing the role of large calcium-based events in place field plasticity in CA1 (Bittner et al. 2015).

Protocol: **In Vivo Patch-Clamp Recording in Awake Head-Fixed Rodents** (Lee and Lee 2016a) describes a detailed training procedure and an efficient patching method for obtaining whole-cell recordings from awake head-fixed animals (Lee et al. 2014).

INTRACELLULAR RECORDINGS IN FREELY MOVING ANIMALS

Although many questions regarding input–output transformations within individual neurons in different behavioral paradigms can be addressed using head-fixed animals, other topics require animals to be non-head-fixed—for example, processing that depends on vestibular input or requires rapid head movements (Monaco et al. 2014). Lee et al. (2006) were the first to demonstrate that intracellular recording (in this case using the whole-cell method) was possible in freely moving animals. These recordings were obtained in anesthetized animals using a head-mounted linear microdrive to advance the pipette electrode; the pipette was subsequently fixed (i.e., “anchored”) in place with dental acrylic and the animal awakened for unrestrained behavior. The recordings could survive severe mechanical disturbances and remain stable for ~20 min while rats explored their environment freely. Other groups have shown that sharp intracellular recordings could be obtained and maintained in freely moving birds (Long et al. 2010; Hamaguchi et al. 2014; Kosche et al. 2015; Vallentin and Long 2015) and mice (English et al. 2014; Schneider et al. 2014) using a head-mountable linear microdrive. The first such recordings were made from bird premotor HVC neurons to measure the subthreshold membrane potential dynamics during singing, and the resulting recordings supported a synaptic chain-like mechanism for birdsong generation (Long et al. 2010). Using this method in freely moving mice, Schneider et al. (2014) demonstrated that the membrane potential of excitatory neurons in auditory cortex depolarizes and decreases its variability immediately before and during bodily movements including locomotion, head movements, and grooming. Additional intracellular recordings in head-fixed mice running in place in concert with optogenetic manipulations revealed that these membrane potential dynamics reflect motor-related corollary discharge originating from secondary motor cortex.

An alternate version of the freely moving whole-cell recording technique uses pipette anchoring combined with a conventional micromanipulator located to the side of the animal (Lee et al. 2009)

instead of a head-mounted microdrive. Conventional micromanipulators not only allow the recording pipette to be held in place more strongly during the anchoring process but also allow the pipette to move horizontally, thus providing a higher probability of obtaining high-quality whole-cell recordings (see Lee and Lee [2016a] for a description of how horizontal movement is used for in vivo patching). This technique has been applied to the study of hippocampal place cells in freely moving rats exploring spatial environments. One such study focused on spikelets, which are all-or-none events with similarities to the action potential but much smaller in amplitude and thus not easily detectable extracellularly. Recordings of hippocampal CA1 pyramidal neurons showed that spikelets occur in the behaving animal and can contribute directly to place cell firing during spatial exploration (Epsztein et al. 2010). In another study, the intracellular features of place cells and silent cells were compared in rats exploring a novel environment. Contrary to expectations from the basic model of neuronal integration, silent cells—unlike place cells—did not display large, spatially tuned inputs at the soma. Furthermore, the excitability of these cells measured even before the rats explored the environment (i.e., for the first time) could partially predict which cells would become place cells, suggesting that intrinsic excitability plays a significant role in place field origin and memory allocation (Epsztein et al. 2011). This method using conventional micromanipulators initially used dental acrylic to fix pipettes in place. However, a limitation of this pipette stabilization technique was that the anchoring process took a relatively long time (~10 min) and thus required that the whole-cell recording be obtained in an initially anesthetized animal that was subsequently awakened for free behavior (Lee et al. 2009). In addition, recording losses during the anchoring process limited the overall success rate of the method (Lee et al. 2009).

To improve this method, the use of dental acrylic was replaced with an ultraviolet (UV)-cured adhesive and UV-transparent collar to anchor the whole-cell pipette in place. This provides highly efficient pipette stabilization with no recording losses. Furthermore, it anchors a pipette in 15 sec, thus allowing the whole-cell recording to be obtained in both anesthetized and awake animals (Lee et al. 2014). This technique has also been applied to the study of cellular mechanisms underlying place cell activity. In particular, injecting a spatially uniform holding current into a silent cell reversibly converted it into a place cell during spatial exploration. Injecting different amounts of current into the silent cell revealed the sudden emergence of large, spatially tuned inputs and place cell spiking above a cell-specific threshold, supporting the idea that nonlinear dendritic mechanisms underlie place cell firing by gating the propagation of spatial inputs to the soma (Lee et al. 2012). These observations were reproduced in drug-free, awake-patched recordings in freely moving animals (Lee et al. 2014).

Protocol: Efficient Method for Whole-Cell Recording in Freely Moving Rodents Using UV-Cured Collar-Based Pipette Stabilization (Lee and Lee 2016b) describes a detailed step-by-step procedure for awake-patched whole-cell recording in freely moving, drug-free rats and mice.

We end this introduction with a comment on whole-cell and sharp intracellular recording methods for freely moving animals. High-quality and long-duration recordings of both types have been obtained successfully. There is, at present, no indication of differences in overall success rates when considering the latest versions of the methods applied to the same brain region (e.g., hippocampal subregion CA1) of the same species (e.g., mice) (English et al. 2014; Lee et al. 2014), although differences might emerge with increased use. However, there are some technical differences inherent to each approach. The ability to achieve low series resistance recordings with the whole-cell configuration makes voltage-clamp experiments possible. On the other hand, multiple recordings can be obtained with the same sharp microelectrode, whereas whole-cell pipettes must be replaced after each recording, as well as each attempt in which negative pressure has been applied.

ACKNOWLEDGMENTS

Work in our laboratories is supported by the Howard Hughes Medical Institute and the Institute for Basic Science, Republic of Korea.

REFERENCES

- Aksay E, Gamkrelidze G, Seung HS, Baker R, Tank DW. 2001. In vivo intracellular recording and perturbation of persistent activity in a neural integrator. *Nat Neurosci* 4: 184–193.
- Bennett C, Arroyo S, Hestrin S. 2013. Subthreshold mechanisms underlying state-dependent modulation of visual responses. *Neuron* 80: 350–357.
- Bittner KC, Grienberger C, Vaidya SP, Milstein AD, Macklin JJ, Suh J, Tonegawa S, Magee JC. 2015. Conjunctive input processing drives feature selectivity in hippocampal CA1 neurons. *Nat Neurosci* 18: 1133–1142.
- Constantinople CM, Bruno RM. 2011. Effects and mechanisms of wakefulness on local cortical networks. *Neuron* 69: 1061–1068.
- Covey E, Kauer JA, Casseday JH. 1996. Whole-cell patch-clamp recording reveals subthreshold sound-evoked postsynaptic currents in the inferior colliculus of awake bats. *J Neurosci* 16: 3009–3018.
- Crochet S, Petersen CCH. 2006. Correlating whisker behavior with membrane potential in barrel cortex of awake mice. *Nat Neurosci* 9: 608–610.
- Domnisoru C, Kinkhabwala AA, Tank DW. 2013. Membrane potential dynamics of grid cells. *Nature* 495: 199–204.
- English DF, Peyrache A, Stark E, Roux L, Vallentin D, Long MA, Buzsáki G. 2014. Excitation and inhibition compete to control spiking during hippocampal ripples: Intracellular study in behaving mice. *J Neurosci* 34: 16509–16517.
- Epsztein J, Lee AK, Chorev E, Brecht M. 2010. Impact of spikelets on hippocampal CA1 pyramidal cell activity during spatial exploration. *Science* 327: 474–477.
- Epsztein J, Brecht M, Lee AK. 2011. Intracellular determinants of hippocampal CA1 place and silent cell activity in a novel environment. *Neuron* 70: 109–120.
- Fee MS. 2000. Active stabilization of electrodes for intracellular recording in awake behaving animals. *Neuron* 27: 461–468.
- Ferster D, Jagadeesh B. 1992. EPSP-IPSP interactions in cat visual cortex studied with in vivo whole-cell patch recording. *J Neurosci* 12: 1262–1274.
- Haider B, Häusser M, Carandini M. 2013. Inhibition dominates sensory responses in the awake cortex. *Nature* 493: 97–100.
- Hamaguchi K, Tschida KA, Yoon I, Donald BR, Mooney R. 2014. Auditory synapses to song premotor neurons are gated off during vocalization in zebra finches. *Elife* 3: e01833.
- Harvey CD, Collman F, Dombeck DA, Tank DW. 2009. Intracellular dynamics of hippocampal place cells during virtual navigation. *Nature* 461: 941–946.
- Kosche G, Vallentin D, Long MA. 2015. Interplay of inhibition and excitation shapes a premotor neural sequence. *J Neurosci* 35: 1217–1227.
- Lee D, Lee AK. 2016a. In vivo patch-clamp recording in awake head-fixed rodents. *Cold Spring Harb Protoc* doi: 10.1101/pdb.prot095802.
- Lee D, Lee AK. 2016b. Efficient method for whole-cell recording in freely moving rodents using UV-cured collar-based pipette stabilization. *Cold Spring Harb Protoc* doi: 10.1101/pdb.prot095810.
- Lee AK, Manns ID, Sakmann B, Brecht M. 2006. Whole-cell recordings in freely moving rats. *Neuron* 51: 399–407.
- Lee AK, Epsztein J, Brecht M. 2009. Head-anchored whole-cell recordings in freely moving rats. *Nat Protoc* 4: 385–392.
- Lee D, Lin B-J, Lee AK. 2012. Hippocampal place fields emerge upon single-cell manipulation of excitability during behavior. *Science* 337: 849–853.
- Lee D, Shtengel G, Osborne JE, Lee AK. 2014. Anesthetized- and awake-patched whole-cell recordings in freely moving rats using UV-cured collar-based electrode stabilization. *Nat Protoc* 9: 2784–2795.
- Long MA, Lee AK. 2012. Intracellular recording in behaving animals. *Curr Opin Neurobiol* 22: 34–44.
- Long MA, Jin DZ, Fee MS. 2010. Support for a synaptic chain model of neuronal sequence generation. *Nature* 468: 394–399.
- Mahon S, Vautrelle N, Pezard L, Slaght SJ, Deniau J-M, Chouvet G, Charpier S. 2006. Distinct patterns of striatal medium spiny neuron activity during the natural sleep–wake cycle. *J Neurosci* 26: 12587–12595.
- Maimon G, Straw AD, Dickinson MH. 2010. Active flight increases the gain of visual motion processing in *Drosophila*. *Nat Neurosci* 13: 393–399.
- Margrie T, Brecht M, Sakmann B. 2002. In vivo, low-resistance, whole-cell recordings from neurons in the anesthetized and awake mammalian brain. *Pflügers Archiv* 444: 491–498.
- Metherate R, Cox CL, Ashe JH. 1992. Cellular bases of neocortical activation: Modulation of neural oscillations by the nucleus basalis and endogenous acetylcholine. *J Neurosci* 12: 4701–4711.
- Monaco JD, Rao G, Roth ED, Knierim JJ. 2014. Attentive scanning behavior drives one-trial potentiation of hippocampal place fields. *Nat Neurosci* 17: 725–731.
- Moore CI, Nelson SB. 1998. Spatio-temporal subthreshold receptive fields in the vibrissa representation of rat primary somatosensory cortex. *J Neurophysiol* 80: 2882–2892.
- Niell CM, Stryker MP. 2010. Modulation of visual responses by behavioral state in mouse visual cortex. *Neuron* 65: 472–479.
- Pei X, Volgushev M, Vidyasagar TR, Creutzfeldt OD. 1991. Whole cell recording and conductance measurements in cat visual cortex in vivo. *NeuroReport* 2: 485–488.
- Polack P-O, Friedman J, Golshani P. 2013. Cellular mechanisms of brain state-dependent gain modulation in visual cortex. *Nat Neurosci* 16: 1331–1339.
- Schmidt-Hieber C, Häusser M. 2013. Cellular mechanisms of spatial navigation in the medial entorhinal cortex. *Nat Neurosci* 16: 325–331.
- Schneider DM, Nelson A, Mooney R. 2014. A synaptic and circuit basis for corollary discharge in the auditory cortex. *Nature* 513: 189–194.
- Steriade M, Timofeev I, Grenier F. 2001. Natural waking and sleep states: A view from inside neocortical neurons. *J Neurophysiol* 85: 1969–1985.
- Tan AYY, Chen Y, Scholl B, Seidemann E, Priebe NJ. 2014. Sensory stimulation shifts visual cortex from synchronous to asynchronous states. *Nature* 509: 226–229.
- Vallentin D, Long MA. 2015. Motor origin of precise synaptic inputs onto forebrain neurons driving a skilled behavior. *J Neurosci* 35: 299–307.
- Wilson RI, Turner GC, Laurent G. 2004. Transformation of olfactory representations in the *Drosophila* antennal lobe. *Science* 303: 366–370.
- Zhu JJ, Connors BW. 1999. Intrinsic firing patterns and whisker-evoked synaptic responses of neurons in the rat barrel cortex. *J Neurophysiol* 81: 1171–1183.

Cold Spring Harbor Protocols

Whole-Cell Recording in the Awake Brain

Doyun Lee and Albert K. Lee

Cold Spring Harb Protoc; doi: 10.1101/pdb.top087304

Email Alerting Service

Receive free email alerts when new articles cite this article - [click here](#).

Subject Categories

Browse articles on similar topics from *Cold Spring Harbor Protocols*.

[Behavioral Assays](#) (99 articles)
[Electrophysiology](#) (104 articles)
[Neuroscience, general](#) (357 articles)
[Patch Clamping](#) (64 articles)

To subscribe to *Cold Spring Harbor Protocols* go to:
<http://cshprotocols.cshlp.org/subscriptions>