Erratum

Cold Spring Harb Protoc 2010; doi: 10.1101/pdb.prot4806

Erratum: Staining for Acetylcholinesterase in Brain Sections

Carol Ann Paul, Barbara Beltz, and Joanne Berger-Sweeney

When this article was first published, the recipe entitled “Sodium acetate buffer (0.2 M, pH 5.0)” (which is used in the acetylcholinesterase stain) listed “sodium acetate” as one of the reagents to be used. However, this reagent should have been “sodium acetate trihydrate,” which will produce a final concentration of 0.2 M sodium acetate in the solution. A corrected version of the recipe is below. In addition, the HTML version of the recipe (doi: 10.1101/pdb.rec11203) has been amended.

Sodium Acetate Buffer (0.2 M, pH 5.0)

Reagent	Quantity (for 2 L)
Sodium acetate trihydrate	54.43 g
Glacial acetic acid	12 mL
H ₂ O	1988 mL
Combine the reagents and adjust the pH to 5.0 with 10 N NaOH.

Cold Spring Harbor Protocols

Erratum: Staining for Acetylcholinesterase in Brain Sections

Carol Ann Paul, Barbara Beltz and Joanne Berger-Sweeney

Cold Spring Harb Protoc; doi: 10.1101/pdb.err101634

Email Alerting Service

Receive free email alerts when new articles cite this article - [click here](#).

Subject Categories

Browse articles on similar topics from *Cold Spring Harbor Protocols*.

To subscribe to *Cold Spring Harbor Protocols* go to:
<http://cshprotocols.cshlp.org/subscriptions>